

# Addressing Controversies in the Management of Ankle Fractures

<sup>1</sup>Amin Kheiran, <sup>2</sup>Jitendra Mangwani

## ABSTRACT

Ankle fractures account for approximately 10% of all fractures and are among the most common orthopedic injuries treated surgically. The incidence of these injuries has increased significantly in the last decade, particularly in the elderly population. Regardless of the method of intervention, the primary goal is restoration of normal anatomy to achieve normal biomechanics, painless function, and prevent long-term posttraumatic degeneration. Surgical treatment carries a potential risk of complications, such as nonunion, implant failure, and soft tissue-related complications. Despite the invention of novel devices, surgical techniques and biomechanical studies for restoration and maintenance of the congruent ankle joint following ankle fractures, several aspects of management of these injuries still remain controversial. The aim of this article is to address these controversies based on the available evidence base.

**Keywords:** Ankle fracture, Posterior malleolus, Stress views of ankle, Syndesmotic injury, Tightrope.

**How to cite this article:** Kheiran A, Mangwani J. Addressing Controversies in the Management of Ankle Fractures. *J Foot Ankle Surg (Asia-Pacific)* 2018;4(3):27-34.

**Source of support:** Nil

**Conflict of interest:** None

## INTRODUCTION

Ankle fractures account for approximately 10% of all fractures and are among the most common orthopedic injuries treated surgically.<sup>1</sup> They have bimodal age distribution with peaks in young males and elderly females.<sup>2</sup> The incidence of these injuries has increased significantly in the last decade, particularly in the elderly population.<sup>3</sup> Ankle fractures should be considered as joint fractures even in the absence of fracture cleft in any of the articular surfaces. The usual mechanism of injury is a rotational force to the ankle. The mechanism of injury is highlighted in different classification systems, including that of Weber<sup>4</sup> and Lauge-Hansen,<sup>5</sup> the two most common

systems in use. It is, however, not entirely clear whether the severity of ankle fractures, based on these classification systems, predicts the outcome of these injuries.<sup>6-8</sup> Regardless of the method of intervention, the primary goal is restoration of normal anatomy to achieve normal biomechanics, painless function, and prevent long-term posttraumatic degeneration. Surgical treatment carries a potential risk of complications, such as nonunion, implant failure, and soft tissue-related complications.<sup>9</sup> These may be caused by surgical factors, fracture pattern, severity of injury, and patient characteristics.<sup>10</sup> The surgical factors include patient selection, timing of surgery, surgical approaches, type of implant, and appropriate rehabilitation program.

Despite the invention of novel devices, surgical techniques and biomechanical studies for restoration and maintenance of the congruent ankle joint following ankle fractures, several aspects of management of these injuries still remain controversial. The aim of this article is to address these controversies based on the available evidence base.

## When is the Best Time to Operate?

Swelling of the skin (Fig. 1) and surrounding soft tissue after ankle fracture can pose a significant challenge to the timing of definitive surgical treatment.<sup>9,10</sup> Avoiding wound complications is of paramount importance and must be considered a high priority.<sup>9</sup> A significant risk



**Fig. 1:** Severely swollen ankle fracture with ecchymosis of the medial side

<sup>1</sup>Specialty Registrar, <sup>2</sup>Consultant

<sup>1,2</sup>Department of Trauma and Orthopaedic Surgery, University Hospitals of Leicester, Leicester, East Midlands, UK

**Corresponding Author:** Jitendra Mangwani, Consultant Department of Trauma and Orthopaedic Surgery, University Hospitals of Leicester, Leicester, East Midlands, UK, Phone: +0044116285085, e-mail: leicesterfoot@icloud.com

**Table 1:** Studies demonstrating the influence of delayed surgery on wound-related complications

Study	Early fixation wound complications	Late fixation wound complications	Recommendation for fixation (time to surgery)
Schepers et al <sup>9</sup>	<24 hrs–4%	>24 hrs–13%	Early
Carragee et al <sup>12</sup>	<24 hrs–5%	>1 week–44%	Early
Hoiness et al <sup>11</sup>	<24 hrs–3%	>24 hrs–18%	Early
Breederveld et al <sup>15</sup>	<24 hrs	5–8 days	Early or late
Konrath et al <sup>16</sup>	<5 days	>5 days	Early or late
Miller et al <sup>17</sup>	<5 days	>5 days–4%	Early or late

of wound-related complications, such as surgical site infection was noted when surgery was delayed 1 week or more.<sup>11</sup> More importantly, this can lead to poor functional outcome.<sup>10</sup> When treated early, an improved quality of anatomical reduction can be achieved.<sup>12</sup> Immediate, definitive surgery is possible only if the soft tissues are not critically injured or extremely vulnerable, usually within first 24 hours after initial trauma.<sup>9,10</sup>

If a delay is contemplated in the definitive management due to excessive swelling, the fracture position should be closely monitored both clinically and radiologically.<sup>13</sup> In cases of trimalleolar or unstable bimalleolar fractures, where a stable reduction cannot be maintained by the plaster, an external fixator should be applied, as loss of reduction will lead to further soft tissue complications.<sup>14</sup> A recent systematic review demonstrated a significant difference in infectious wound complications for patients who underwent surgery after a delay for a closed ankle fracture.<sup>9,11,12,15–17</sup> Published data regarding the influence of delayed surgery on the outcome or soft tissue-related complication are limited (Table 1).

## Fracture Blisters

The incidence of fracture blisters in ankle fractures has been reported to be as high as 7%.<sup>18,19</sup> Blisters have significant impact on decision-making in both nonoperative and operative management. Fracture blisters are thought to be the result of a cleavage injury at the junction of dermis and epidermis. Such superficial shearing injuries are called fracture blisters.<sup>20</sup> Anatomical areas, such as the ankle, with poor muscle and adipose tissue cover are especially prone.<sup>18</sup> These blisters can appear as clear or hemorrhagic (sanguineous) blisters (Fig. 2). Clear blisters lie completely within the epidermis, whereas the hemorrhagic type often extends deeper into the dermis, compromising the crossing microcirculation.<sup>21</sup> While some studies support the view that allowing the blisters to resolve before any surgical intervention is desirable,<sup>18,19</sup> others have validated a treatment protocol for unroofing the blister surface and application of silver sulfadiazine antibiotic cream until the swelling of skin permits surgery


**Fig. 2:** Hemorrhagic fracture blisters in a patient with a high-energy injury closed ankle fracture

and the blister appeared re-epithelialized, on average after 7 days.<sup>22</sup> There is no clear consensus on how best to manage ankle fractures with associated blisters. The presence of blisters, particularly the hemorrhagic ones, indicates significant injury to the soft tissues and alternative strategies, such as different surgical approaches, use of minimally invasive techniques, or staged fixation to achieve as near anatomic reduction of the ankle mortise should be adopted to allow adequate resuscitation of traumatized soft tissue envelope.

## Stability in Isolated Lateral Malleolar Fractures

Almost 70% of ankle fractures are stable.<sup>3</sup> Stable ankle fractures will not displace on physiological axial loading when deep deltoid ligament is functionally intact, by providing checkrein and maintaining further stability.<sup>23</sup> A 1-mm lateral talar shift is known to decrease the contact area between talus and tibia by 42%. Incongruity of joint surfaces predisposes to an irreversible condition, such as posttraumatic osteoarthritis.<sup>24</sup> Historically, any fibular displacement was thought to cause talar displacement or shift.<sup>25</sup> Two studies have shown that in isolated lateral malleolar fractures, a functionally intact deep deltoid ligament acts as checkrein and prevents lateral talar or mortise displacement, providing further stability and normal ankle anatomy even during weight-bearing.<sup>26,27</sup> They also stated that apparent fibular displacement is often misleading.


Current practice relies upon different clinical and radiological methods to identify stable isolated lateral malleolar fractures and help dictate an appropriate treatment strategy. These are categorized based on clinical or radiological findings. Medial tenderness, ecchymosis, or swelling has been used clinically to delineate a potential injury of the deltoid ligament, suggesting an unstable morphology.<sup>28,29</sup> A systematic review of the literature



**Fig. 3:** Gravity stress test; positioning of a patient and C-arm "X-ray" machine



**Fig. 4:** Manual external rotation test



**Fig. 5:** Anteroposterior X-rays demonstrating Weber B fracture of the ankle and widening of medial clear space of > 5 mm indicating deltoid ligament injury (positive gravity stress test)

looking at diagnostic modalities to assess the integrity of the deltoid ligament in adult ankle fractures (supination external rotation injuries) concluded that clinical signs, such as swelling and ecchymosis on medial side with associated tenderness, initial radiographic findings, and the Lauge–Hansen classification systems are poor predictors of deltoid ligament injury and ankle stability.<sup>30</sup>

Stress X-ray is considered the gold standard to identify stable or unstable ankle fractures (Figs 3 to 5). Manual and the gravity external rotation tests are two most common ways to perform stress X-rays.<sup>66</sup> Medial clear space widening of more than 5 mm is regarded as a reliable indicator of unstable ankle fracture. Both tests have proven to be effective.<sup>31</sup> However, the amount of force applied when performing an external rotation stress radiograph (dynamic test) is not well defined and purely determined by the patient's pain level and assessor's familiarity with the procedure. Manual stress test requires more time and radiation exposure, whereas gravity stress test necessitates X-ray education.

### Does an Abnormal Stress Test equate to Surgery?


Based on magnetic resonance imaging (MRI) studies, 90% of patients with positive stress radiographs have partial tear of deep deltoid ligament. These fractures can heal without surgical intervention as long as they are reduced adequately and immobilized.<sup>31,32</sup> Stress radiographs can overestimate the need for surgery.<sup>33,34</sup> If in doubt, clinician is suggested to obtain standing radiographs to differentiate between stable and unstable ankle.<sup>35</sup> About 89 to 100% of patients with abnormal stress X-rays were eventually stable on standing radiographs.<sup>33,35</sup>

The indication for surgery should not be based on the absolute value of one parameter, but on the combination of several measures. Close monitoring/follow-up is essential if nonoperative treatment is chosen despite a positive stress test, because subluxation or displacement of the ankle joint is still possible. The MRI can be useful in individual cases.<sup>33</sup>

### Posterior Malleolus Fractures: Need for Preoperative Computed Tomography and Which Approach to fix?

The prevalence of posterior malleolar injury in ankle fractures has been reported to be as high as 44%. The functional outcome following ankle fractures with posterior malleolar fragment is often not satisfactory,<sup>36</sup> particularly when associated with syndesmotic injury. Operative management continues to be controversial. Morphology of fracture has received far less consideration in common fracture classification systems and treatment algorithms. The fracture lines associated with posterior malleolar fractures are variable.<sup>36</sup> The common fracture patterns are categorized into three types: (1) The posterolateral-oblique type (67%), (2) the medial-extension type (19%), and (3) the





**Figs 6A and B:** Postoperative lateral X-rays demonstrating fixation of posterior malleolus fracture using (a) anteroposterior cannulated screw, and (b) plate and screws through posterior approach

small-shell type (14%).<sup>36</sup> Knowledge of this pathoanatomy and careful scrutiny of the preoperative imaging are essential for approaching these fractures. Preoperative computed tomography (CT) is useful to delineate anatomy of the fracture, presence of comminution, impaction of the fragment, and to plan the approach.<sup>36,37</sup>

Traditionally, the decision to fix posterior malleolus has been based upon the amount of articular surface involved. Other factors to consider are a posterior subluxation of the talus, an articular step-off of more than 2 mm, instability after fibular fixation, or residual syndesmotom widening or malreduced mortise. A prospective study with long-term follow-up period has demonstrated fair-to-good outcomes when fracture fragment is involved less than 25% of articular surfaces and managed nonoperatively.<sup>38</sup>

Two most common ways to fix posterior malleolus fixation are anteroposterior (AP) or posteroanterior. There is controversy with regard to screw fixation *vs* plating (Fig. 6) via posterior approach.<sup>38,39</sup>

It has been shown that when posterior malleolus is fractured, posterior syndesmotom ligaments are intact and attached to the fragment.<sup>39</sup> In a cadaveric study assessing the biomechanics of syndesmosis after internal fixation of the posterior malleolus, 70% of syndesmotom stability could be established in contrast to restoration of 40% by fixing the syndesmosis alone.<sup>39</sup> It is important to ensure that posterior fragment is well reduced and there is no subluxation of the ankle joint once medial and/or lateral malleoli are stabilized.<sup>40</sup> A prospective study has shown a significant difference in outcomes comparing patients with unstable ankle fractures associated with or without posterior malleolus fracture (fixed or not fixed). The presence of posterior malleolus fracture indicates high-energy trauma and seems to result in worse outcomes at 1 year after intervention.<sup>41</sup>



**Fig. 7:** Mortise view radiograph of an ankle fracture with syndesmotom disruption


### Syndesmotom Injuries: To fix or Not to fix?

Over 90% of the total resistance to lateral displacement of the fibula is provided by the three syndesmotom ligaments, and injury to one or more of them results in weakening, abnormal movement of the joint, and instability (Fig. 7). Although many mechanisms for syndesmotom injury have been reported, the most common is external rotation of the foot and, to a lesser extent, forced dorsiflexion of the ankle with axial loading. In most complete syndesmotom disruptions, external rotation causes a Weber C or Weber B fracture with widening of the mortise and, occasionally, a Maisonneuve fracture.<sup>42</sup>

In up to 13% of all ankle fractures, and in 20% of patients requiring internal fixation, there will be an associated injury to the syndesmosis. These injuries can create a diagnostic challenge and there is a lack of consensus on optimal method of treatment. There are controversies with regard to the type of fixation device (screw *vs* TightRope<sup>®</sup>), characteristics and position of the screw, the type of cortical fixation, number of screws, and whether the screw should be retained or removed prior to weight-bearing. It remains unclear whether these technical aspects of surgery affect the clinical outcome.<sup>43</sup>

### How to assess Syndesmotom Injuries?

Tibiofibular clear space and overlap (Fig. 8) are used as a common radiological assessment when suspecting syndesmosis disruption. A normal tibiofibular clear space is defined as a distance between the lateral border of the posterior tubercle and the medial border of the fibula.<sup>42</sup> The tibiofibular overlap is the distance between the medial border of the fibula and the lateral border of the anterior distal tibial tubercle.<sup>42</sup> These are measured at 1 cm proximal to the ankle joint.<sup>44</sup> A normal tibiofibular clear space should be less than 6 mm on both AP or



**Fig. 8:** Mortise view X-ray of ankle illustrating radiological parameters for assessment of syndesmosis: (A and B) tibiofibular overlap ( $N > 6$  mm); (B and C) tibiofibular clear space ( $N < 6$  mm); and (D and E) medial clear space ( $N < 5$  mm)

mortise views, whereas tibiofibular overlap of  $> 6$  mm on the AP view or  $> 1$  mm on the mortise view suggest an intact syndesmosis.<sup>42,44</sup> However, these indices for confirmation of syndesmosis disruption could not detect externally rotated malreduction of syndesmosis of up to  $30^\circ$ .<sup>45</sup> The position of the ankle greatly influenced these measurements, and some authors believe there are no optimal radiological parameters to assess the integrity of the syndesmosis.<sup>46-48</sup> Although CT scan has been considered more sensitive than plain radiographs to detect syndesmotic injuries, MRI has become the modality of choice to delineate the syndesmotic integrity.<sup>49</sup> However, MRI has not been considered as a routine modality of investigation due to its cost-implications.<sup>50</sup>

### How to assess Syndesmotic Stability during Operation?

Many orthopedic surgeons evaluate the need for syndesmotic fixation intraoperatively by pulling laterally on the fibula with a bone hook (Fig. 9). Widening of the syndesmosis by more than 2 mm on the mortise view suggests the need for fixation. Despite being a popular diagnostic tool, the “hook test” is poorly described in the literature and can be difficult to interpret.<sup>51</sup> Candal-Couto et al<sup>52</sup> assessed the reliability of this test in a cadaver model by sequentially dividing the ligaments of the syndesmosis and finally the deltoid ligament. They showed that the AP and mortise views correlated poorly with the observed clinical syndesmotic injury.<sup>52</sup> However, performing the hook test in the sagittal plane (the sagittal-shift test) appeared to be a more sensitive assessment of inferior tibiofibular instability.<sup>52</sup> Similarly, fluoroscopic examination following the application of an external rotation stress has been shown to demonstrate



**Fig. 9:** Anteroposterior X-ray of ankle demonstrating assessment of syndesmosis intra operatively using “Hook test”

syndesmotic instability.<sup>53</sup> However, there is no consensus on how much force is needed to accurately identify the potential pathology.<sup>42</sup> Some authors suggest that arthroscopy is required for the accurate diagnosis of syndesmotic disruption.<sup>49</sup> Damage to the tibiofibular syndesmosis can be diagnosed accurately in 100% of cases by arthroscopy of the ankle, compared with only 48% by AP radiography, 64% by mortise views, and 96% with MRI.<sup>43,50</sup>

### Which Device to use to fix Syndesmotic Injuries?

Most surgeons advocate the use of metal screws for stabilization of the syndesmosis, but opinions vary with regard to the characteristics of the procedure or type of the device.<sup>42</sup>

Randomized studies comparing metal and bioabsorbable screws have demonstrated that both techniques are equally effective in fixation of a syndesmotic disruption, with patients more likely to return to their previous level of activity when treated with a bioabsorbable rather than a metal screw.<sup>54</sup> However, concerns about the use of bioabsorbable materials include osteolysis, foreign body reaction, late inflammatory reaction, and osteoarthritis due to polymer debris entering the joint.<sup>55,56</sup> TightRopes<sup>®</sup> may be used with placement of a heavy suture, which is looped and tightened through cortical button anchors on either side of the ankle.<sup>57</sup> TightRope<sup>®</sup> has shown similar outcome, but quicker time to recovery or return to work based on a systematic review of the literature.<sup>58</sup> A recent randomized controlled trial (RCT) has concluded that dynamic fixation with TightRope<sup>®</sup> appears to result in better functional and radiological outcomes in acute ankle syndesmotic rupture. They also demonstrated that TightRope<sup>®</sup> offers good stabilization without failure or loss of reduction and subsequently the reoperation rate was significantly lower than the conventional metallic screw fixation.<sup>59</sup>

Regardless of the method of fixation, patients who required syndesmotic fixation in addition to their malleolar fracture stabilization showed poorer outcome at 12 months. This information is important for patient counseling and managing their expectation regarding recovery and regaining function after injury.<sup>40</sup>

### Retain or remove the Syndesmotic Screw?

There is no consensus whether syndesmotic screw should be removed prior to weight-bearing or left in place indefinitely. Fixation with a screw provides rigid fixation of the distal tibiofibular joint where physiological micromovement has been shown to occur.<sup>47,60</sup> Therefore, leaving it in place may contribute to abnormal ankle movement, which, in turn, may result in loosening or fatigue fracture of the screw.<sup>61,62</sup>

In a literature review conducted by Schepers et al,<sup>43</sup> limited level I studies were available on the absolute requirement for removal of the syndesmotic screw. Most included studies found no difference in functional outcome between retained and removed metalwork. Removal of syndesmotic screws is usually not performed before 4 to 6 months.

A comparative study by Stuart and Panchbhavi,<sup>63</sup> evaluating 137 syndesmotic fixation using 3.5 *vs* 4.5 mm screws, demonstrated no difference in loss of reduction, but 3.5-mm screws were more likely to break.

### Postoperative Rehabilitation: Early Mobilization vs Plaster Cast

In a literature review of 31 RCTs concerning rehabilitation of ankle fractures, common complications elucidated included pain, stiffness, weakness, and swelling.<sup>64</sup> All these are recognized as barriers to overcome for successful rehabilitation. Evidence is lacking regarding intervention following conservative management, with more evidence available on interventions following surgery.<sup>13</sup> A prospective RCT recruiting 100 patients after ankle surgery (open reduction and internal fixation) compared an immobilizing cast with a functional ankle brace. Results showed that functional outcome was similar at 2 years follow-up although brace carried higher risk of wound complications.<sup>65</sup>

A combination of early mobilization, early commencement of weight-bearing, and the use of a removable immobilization device, in conjunction with exercise showed a positive effect on ankle range of motion.<sup>64</sup> A systematic review identified an increased risk of wound complications when ankle was mobilized early, but patients returned to activity or work quicker. It was observed that patients' compliance is a predisposing factor along with skin condition or other comorbidities

(e.g., peripheral vascular disease, diabetes). It is important to consider patient factors, particularly their ability to correctly apply and use a temporary immobilization device and their compliance with directed exercise regimes, as these variables can influence overall effectiveness of the intervention.<sup>13,64,65</sup>

### CONCLUSION

Although ankle fracture is a common injury, there are still controversies with regard to its optimum management. It is imperative to understand basic biomechanics, pattern, and mechanism of injury. Early intervention to achieve anatomical reduction and stabilize the fracture is recommended. If possible, the fractures should preferably be fixed definitively within the first 24 hours. A delay of more than 1 week gives a significant rise in surgical-related complication, particularly wound infections, which lowers functional outcome and patient satisfaction. Diagnostic tests and management of syndesmotic injuries are still controversial, but sagittal plane instability should be recognized in addition to coronal plane. Associated syndesmotic injury and posterior malleolus fracture lead to poor functional outcome and patients should be counseled appropriately. Early mobilization after primary surgery leads to quicker recovery and early return to work, but may carry higher risk of wound-related complications.

### REFERENCES

1. Court-Brown CM, Caesar B. Epidemiology of adult fractures: a review. *Injury* 2006 Aug;37(8):691-697.
2. Court-Brown C, McBirnie J, Wilson G. Adult ankle fractures: an increasing problem? *Acta Orthop Scand* 1998 Feb;69(1):43-47.
3. Kannus P, Palvanen M, Niemi S, Parkkari J, Järvinen M. Increasing number and incidence of low-trauma ankle fractures in elderly people: Finnish statistics during 1970-2000 and projections for the future. *Bone* 2002 Sep;31(3):430-433.
4. Weber, BG. *Die verletzungen des oberen sprunggelenkes*. 2nd ed. Berne: Verlag Hans Huber; 1972.
5. Lauge-Hansen N. Fractures of the ankle. III. Genetic roentgenologic diagnosis of fractures of the ankle. *Am J Roentgenol Radium Ther Nucl Med* 1954 Mar;71(3):456-471.
6. Broos PL, Bisschop AP. Operative treatment of ankle fractures in adults: correlation between types of fracture and final results. *Injury* 1991 Sep;22(5):403-406.
7. Egol KA, Tejwani NC, Walsh MG, Capla EL, Koval KJ. Predictors of short-term functional outcome following ankle fracture surgery. *J Bone Joint Surg Am* 2006 May;88(5):974-979.
8. Shah NH, Sundaram RO, Velusamy A, Braithwaite IJ. Five year functional outcome analysis of ankle fracture fixation. *Injury* 2007 Nov;38(11):1308-1312.
9. Schepers T, Lieshout EM, De Vries MR, Van der Elst M. Increased rates of wound complications with locking



- plates in distal fibular fractures. *Injury* 2013 Oct;42(10):1125-1129.
10. SooHoo NF, Krenek L, Eagan MJ, Gurbani B, Ko CY, Zingmond DS. Complication rates following open reduction and internal fixation of ankle fractures. *J Bone Joint Surg Am* 2009 May;91(5):1042-1049.
  11. Höiness P, Engebretsen L, Strömsöe K. The influence of perioperative soft tissue complications on the clinical outcome in surgically treated ankle fractures. *Foot Ankle Int* 2001 Aug;22(8):642-648.
  12. Carragee EJ, Csongradi JJ. Increased rates of complications in patients with severe ankle fractures following interinstitutional transfers. *J Trauma* 1993 Nov;35(5):767-771.
  13. Mehta SS, Rees K, Cutler L, Mangwani J. Understanding risks and complications in the management of ankle fractures. *Indian J Orthop* 2014 Sep;48(5):445-452.
  14. Rammelt S, Endres T, Grass R, Zwipp H. The role of external fixation in acute ankle trauma. *Foot Ankle Clin* 2004 Sep;9(3):455-474.
  15. Breederveld RS, van Straaten J, Patka P, van Mourik JC. Immediate or delayed operative treatment of fractures of the ankle. *Injury* 1998 Nov;19(6):436-438.
  16. Konrath G, Karges D, Watson JT, Moed BR, Cramer K. Early versus delayed treatment of severe ankle fractures: a comparison of results. *J Orthop Trauma* 1995 Oct;9(5):377-380.
  17. Miller AG, Margules A, Raikin SM. Risk factors for wound complications after ankle fracture surgery. *J Bone Joint Surg Am* 2012 Nov;94(22):2047-2052.
  18. Uebbing CM, Walsh M, Miller JB, Abraham M, Arnold C. Fracture blisters. *West J Emerg Med* 2011 Feb;12(1):131-133.
  19. Varela CD, Vaughan TK, Carr JB, Slemmons BK. Fracture blisters: clinical and pathological aspects. *J Orthop Trauma* 1993 Oct;7(5):417-427.
  20. Morel-Lavallée M. Degloving of skin and underlying tissues. *Arch Gen Med* 1863;1:20-38, 172-200, 300-332.
  21. Giordano CP, Koval KJ. Treatment of fracture blisters: a prospective study of 53 cases. *J Orthop Trauma* 1995 Apr;9(2):171-176.
  22. Strauss EJ, Petrucelli G, Bong M, Koval KJ, Egol KA. Blisters associated with lower-extremity fracture: results of a prospective treatment protocol. *J Orthop Trauma* 2006 Oct;20(9):618-622.
  23. Michelson JD, Hamel AJ, Buczek FL, Sharkey NA. Kinematic behavior of the ankle following malleolar fracture repair in a high-fidelity cadaver model. *J Bone Joint Surg Am* 2002 Nov;84-A(11):2029-2038.
  24. Ramsey PL, Hamilton W. Changes in tibiotalar area of contact caused by lateral talar shift. *J Bone Joint Surg Am* 1976 Apr;58(3):356-357.
  25. Yablon I G, Heller F G, Shouse L. The key role of the lateral malleolus in displaced fractures of the ankle. *J Bone Joint Surg Am* 1977 Mar;59(2):169-173.
  26. Michelson JD, Magid D, Ney DR, Fishman EK. Examination of the pathologic anatomy of ankle fractures. *J Trauma* 1992 Jan;32(1):65-70.
  27. Harper MC. The short oblique fracture of the distal fibula without medial injury: an assessment of displacement. *Foot Ankle Int* 1995 Apr;16(4):181-186.
  28. DeAngelis NA, Eskander MS, French BG. Does medial tenderness predict deep deltoid ligament incompetence in supination-external rotation type ankle fractures? *J Orthop Trauma* 2007 Apr;21(4):244-247.
  29. Pettrone FA, Gail M, Pee D, Fitzpatrick T, Van Herpe LB. Quantitative criteria for prediction of the results after displaced fracture of the ankle. *J Bone Joint Surg Am* 1983 Jun;65(5):667-677.
  30. van den Bekerom MP, Mutsaerts EL, van Dijk CN. Evaluation of the integrity of the deltoid ligament in supination external rotation ankle fractures: a systematic review of the literature. *Arch Orthop Trauma Surg* 2009 Feb;129(2):227-235.
  31. Koval KJ, Egol KA, Cheung Y, Goodwin DW, Spratt KF. Does a positive ankle stress test indicate the need for operative treatment after lateral malleolus fracture? A preliminary report. *J Orthop Trauma* 2007 Aug;21(7):449-455.
  32. Cheung Y, Perrich KD, Gui J, Koval KJ, Goodwin DW. MRI of isolated distal fibular fractures with widened medial clear space on stressed radiographs: which ligaments are interrupted? *AJR Am J Roentgenol* 2009 Jan;192(1):W7-W12.
  33. Weber M, Burmeister H, Flueckiger G, Krause FG. The use of weightbearing radiographs to assess the stability of supination-external rotation fractures of the ankle. *Arch Orthop Trauma Surg* 2010 May;130(5):693-698.
  34. Egol KA, Amirtharajah M, Tejwani NC, Capla EL, Koval KJ. Ankle stress test for predicting the need for surgical fixation of isolated fibular fractures. *J Bone Joint Surg Am* 2004 Nov;86-A(11):2393-2398.
  35. Hoshino CM, Nomoto EK, Norheim EP, Harris TG. Correlation of weightbearing radiographs and stability of stress positive ankle fractures. *Foot Ankle Int* 2012 Feb;33(2):92-98.
  36. Haraguchi N, Haruyama H, Toga H, Kato F. Pathoanatomy of posterior malleolar fractures of the ankle. *J Bone Joint Surg Am* 2006 May;88(5):1085-1092.
  37. Büchler L, Tannast M, Bonel HM, Weber M. Reliability of radiologic assessment of the fracture anatomy at the posterior tibial plafond in malleolar fractures. *J Orthop Trauma* 2009 Mar;23(3):208-212.
  38. De Vries JS, Wiggins AJ, Sierevelt IN, Schaap GR. Long term results of ankle fractures with a posterior malleolar fragment. *J Foot Ankle Surg* 2005 May-Jun;44(3):211-217.
  39. Gardner MJ, Brodsky A, Briggs SM, Nielson JH, Lorich DG. Fixation of posterior malleolar fractures provides greater syndesmotic stability. *Clin Orthop Relat Res* 2006 Jun;447:165-171.
  40. Egol KA, Pahk B, Walsh M, Tejwani NC, Davidovitch RI, Koval KJ. Outcome after stable ankle fracture: effect of syndesmotic stabilization. *J Orthop Trauma* 2010 Jan;24(1):7-11.
  41. Tejwani NC, Pahk B, Egol KA. Effect of posterior malleolus fracture on outcome after unstable ankle fracture. *J Trauma* 2010 Sep;69(3):666-669.
  42. Dattani R, Patnaik S, Kantak A, Srikanth B, Selvan TP. Injuries to the tibiofibular syndesmosis. *J Bone Joint Surg Br* 2008 Apr;90(4):405-410.
  43. Schepers T. To retain or remove the syndesmotic screw: a review of literature. *Arch Orthop Trauma Surg* 2011 Jul;131(7):879-883.
  44. Xenos JS, Hopkinson WJ, Mulligan ME, Olson EJ, Popovic NA. The tibiofibular syndesmosis: evaluation of the ligamentous structures, methods of fixation, and radiographic assessment. *J Bone Joint Surg Am* 1995 Jun;77(6):847-856.
  45. Marmor M, Hansen E, Han HK, Buckley J, Matityahu A. Limitations of standard fluoroscopy in detecting rotational malreduction of the syndesmosis in an ankle fracture model. *Foot Ankle Int* 2011 Jun;32(6):616-622.
  46. Beumer A, van Hemert WL, Niesing R, Entius CA, Ginai AZ, Mulder PG, Swierstra BA. Radiographic measurement of the

- distal tibiofibular syndesmosis has limited use. *Clin Orthop* 2004 Jun;423:227-234.
47. Beumer A, Valstar ER, Garling EH, Niesing R, Heijboer RP, Ranstam J, Swierstra BA. Kinematics before and after reconstruction of the anterior syndesmosis of the ankle: a prospective radiostereometric and clinical study in five patients. *Acta Orthop* 2005 Oct;76(5):713-720.
48. Schuberth JM, Collman DR, Rush SM, Ford LA. Deltoid ligament integrity in lateral malleolar fractures: a comparative analysis of arthroscopic and radiographic assessments. *J Foot Ankle Surg* 2004 Jan-Feb;43(1):20-29.
49. Takao M, Ochi M, Naito K, Iwata A, Kawasaki K, Tobita M, Miyamoto W, Oae K. Arthroscopic diagnosis of tibiofibular syndesmosis disruption. *Arthroscopy* 2001 Oct;17(8):836-843.
50. Takao M, Ochi M, Oae K, Naito K, Uchio Y. Diagnosis of a tear of the tibiofibular syndesmosis: the role of arthroscopy of the ankle. *J Bone Joint Surg Br* 2003 Apr;85(3):324-329.
51. van den Bekerom MP, Hogervorst M, Bolhuis HW, van Dijk CN. Operative aspects of the syndesmotic screw: review of current concepts. *Injury* 2008 Apr;39(4):491-498.
52. Candal-Couto JJ, Burrow D, Bromage S, Briggs PJ. Instability of the tibiofibular syndesmosis: have we been pulling in the wrong direction? *Injury* 2004 Aug;35(8):814-818.
53. Jenkinson RJ, Sanders DW, Macleod MD, Domonkos A, Lydestadt J. Intraoperative diagnosis of syndesmosis injuries in external rotation ankle fractures. *J Orthop Trauma* 2005 Oct;19(9):604-609.
54. Sinisaari IP, Luthje PM, Mikkonen RH. Ruptured tibio-fibular syndesmosis: comparison study of metallic to bioabsorbable fixation. *Foot Ankle Int* 2002 Aug;23(8):744-748.
55. Thordarson IP, Samuelson M, Shepherd LE, Merkle PF, Lee J. Bioabsorbable versus stainless steel screw fixation of the syndesmosis in pronation-lateral rotation ankle fractures: a prospective randomized trial. *Foot Ankle Int* 2001 Apr;22(4):335-338.
56. Kaukonen JP, Lamberg T, Korkala O, Pajarinen J. Fixation of syndesmotic ruptures in 38 patients with a malleolar fracture: a randomized study comparing a metallic and a bioabsorbable screw. *J Orthop Trauma* 2005 Jul;19(6):392-395.
57. Thornes B, Walsh A, Hislop M, Murray P, O'Brien M. Suture-endobutton fixation of ankle tibio-fibular diastasis: a cadaver study. *Foot Ankle Int* 2003 Feb;24(2):142-146.
58. Schepers T. Acute distal tibiofibular syndesmosis injury: a systematic review of suture-button versus syndesmotic screw repair. *Int Orthop* 2012 Jun;36(6):1199-1206.
59. Laflamme M, Belzile EL, Bédard L, van den Bekerom MP, Glazebrook M, Pelet S. A prospective randomized multicenter trial comparing clinical outcomes of patients treated surgically with a static or dynamic implant for acute ankle syndesmosis rupture. *J Orthop Trauma* 2015 May;29(5):216-223.
60. Beumer A, Valstar ER, Garling EH, Niesing R, Ranstam J, Löfvenberg R, Swierstra BA. Kinematics of the distal tibiofibular syndesmosis: radiostereometry in 11 normal ankles. *Acta Orthop Scand* 2003 Jun;74(3):337-343.
61. Needleman RL, Skrade DA, Stiehl JB. Effect of the syndesmotic screw on ankle motion. *Foot Ankle* 1989 Aug;10(1):17-24.
62. Heim D, Heim U, Regazzoni P. Malleolar fractures with ankle joint instability: experience with the positioning screw. *Unfallchirurgie* 1993 Oct;19(5):307-312.
63. Stuart K, Panchbhavi VK. The fate of syndesmotic screws. *Foot Ankle Int* 2011 May;32(5):S519-S525.
64. Lin CW, Donkers NA, Refshauge KM, Beckenkamp PR, Khera K, Moseley AM. Rehabilitation for ankle fractures in adults. *Cochrane Database Syst Rev* 2012 Nov;11:CD005595.
65. Lehtonen H, Järvinen TL, Honkonen S, Nyman M, Vihtonen K, Järvinen M. Use of a cast compared with a functional ankle brace after operative treatment of an ankle fracture. A prospective, randomized study. *J Bone Joint Surg Am* 2003 Feb;85-A(2):205-211.
66. Schock HJ, Pinzur M, Manion L, Stover M. The use of gravity or manual-stress radiographs in the assessment of supination-external rotation fractures of the ankle. *J Bone Joint Surg Br* 2007 Aug;89(8):1055-1059.